*Journal of Immunological Methods*, 152 (1992) 269–270

© 1992 Elsevier Science Publishers B.V. All rights reserved 0022-1759/92/$05.00

# Contents to volume 152 (1992)

**Number 1 – 31 July 1992**

*Research reports*

Parameters affecting the adsorption of ligands to polyvinyl chloride plates in enzyme immunoassays
*F.E. Zalazar, G.A. Chiabrando, M.A.J. Aldao and M.A. Vides (Argentina)* 1

Increased ELISA sensitivity using a modified method for conjugating horseradish peroxidase to monoclonal antibodies
*S. Madersbacher, H. Wolf, R. Gerth and P. Berger (Austria)* 9

Generation of monoclonal antibodies to human lymphocyte cell surface antigens using insect cells expressing recombinant proteins
*M. De Boer, L. Conroy, H.Y. Min and J. Kwekkeboom (USA, Netherlands)* 15

Evaluation of a flow cytometric model for monitoring HIV antigen expression in vitro
*C.A. Heynen and T.J. Holzer (USA)* 25

Detection of mycoplasma contamination through modulation (stimulation or inhibition) of thymidine incorporation by unstimulated mouse spleen cells
*R. Alzani, R. Rossi, E. Cozzi, D. Trizio and F. Marcucci (Italy)* 35

Chimaeric protein A/protein G and protein G/alkaline phosphatase as reporter molecules
*S. Sun and A.M. Lew (China, Australia)* 43

Agonistic properties of anti-B cell antibodies purified on staphylococcal protein A may be due to contaminating protein A
*E. Jakobson, B. Axelsson and S. Paulie (Sweden)* 49

Determination of monoclonal antibody specificity by immunoadsorption and Western blotting
*R. Schuh, E. Kremmer, E. Ego, M. Wasiliu and S. Thierfelder (Germany)* 59

A new approach to the generation of human or murine antibody producing hybridomas
*P.G.A. Steenbakkers, F.C.M. Van Meel and W. Olijve (Netherlands)* 69

Production of monoclonal antibodies against the human anaphylatoxin C5a des Arg and their application in the neoepitope-specific sandwich-ELISA for the quantification of C5a des Arg in plasma
*K. Bergh and O.-J. Iversen (Norway)* 79

Novel vectors for the expression of antibody molecules using variable regions generated by polymerase chain reaction
*M.J. Coloma, A. Hastings, L.A. Wims and S.L. Morrison (USA)* 89

Comparison of Freund's and Ribi adjuvants for inducing antibodies to the synthetic antigen (TG)-AL in rabbits
*B.J. Deeb, R.F. DiGiacomo, L.L. Kunz and J.L. Stewart (USA)* 105

Comparison of methodologies to measure human lung histamine
*R.A. Erger and T.B. Casale (USA)* 115

Establishment of stable $CD8^+$ suppressor T cell clones and the analysis of their suppressive function
*F.-y. Hu, Y. Asano, K. Sano, T. Inoue, M. Furutani-Seiki and T. Tada (Japan)* 123

Detection of *Salmonella enteritidis* in environmental samples by monoclonal antibody-based ELISA
*R.L. Brigmon, S.G. Zam, G. Bitton and S.R. Farrah (USA)* 135

*Book review* 143

*Notes to authors* 145

**Number 2 – 10 August 1992**

*Immunological standardization* 147

*Research reports*

Isolation of sequences from a random-sequence expression library that mimic viral epitopes
*J.A. Lenstra, J.H.F. Erkens, J.G.A. Langeveld, W.P.A. Posthumus, R.H. Meloen, F. Gebauer, I. Correa, L. Enjuanes and K.K. Stanley (Netherlands, Spain, Germany)* 149

Kinetics of killing by monoclonal cytotoxic T lymphocytes. I. Colorimetric detection of cryptic CTL determinants on adherent target cells and survivorship analysis
*J.R. Rodgers, S.M. Shawar, M.M. Guenther and R.R. Rich (USA)* 159

A flow-cytometric method to study DNA fragmentation in lymphocytes
*F. Ojeda, M.I. Guarda, C. Maldonado and H. Folch (Chile)* 171

Biological activities of polyethylene-glycol immunoglobulin conjugates. Resistance to enzymatic degradation
*C. Cunningham-Rundles, Z. Zhuo, B. Griffith and J. Keenan (USA)* 177

Characterization of monoclonal antibodies physically adsorbed onto polystyrene latex particles
*R. Van Erp, Y.E.M. Linders, A.P.G. Van Sommeren and T.C.J. Gribnau (Netherlands)* 191

U937 cells stimulated with opsonised zymozan particles provide a convenient laboratory source of tumour necrosis factor $\alpha$
*W.G. Jiang, M.C.A. Puntis and M.B. Hallett (UK)* 201

Evaluation of biological response modifiers in the enhancement of tumor uptake of technetium-99m labeled macromolecules. A preliminary report
*M.L. Thakur, J. DeFulvio, J. Tong, E. John, M.R. McDevitt and I. Damjanov (USA)* 209

Analysis of B lymphocyte cross-reactivity at the single cell level
*D.M. Klinman (USA)* 217

Quantitative dot-blot assay for proteins using enhanced chemiluminescence
*E. Heinicke, U. Kumar and D.G. Munoz (Canada)* 227

Class I restricted CTL recognition of a soluble protein delivered by liposomes containing lipophilic polylysines
*S. Nair, X. Zhou, L. Huang and B.T. Rouse (USA)* 237

Differentiation-related changes in quantitative binding of immunomagnetic beads
*I.D. Swann, G.B. Dealtry and D. Rickwood (UK)* 245

Isolation of functionally active intraepithelial lymphocytes and enterocytes from human small and large intestine
*C. Lundqvist, M.-L. Hammarström, L. Athlin and S. Hammarström (Sweden)* 253

*Book reviews* 265

*Announcement from the publisher* 267

*Contents to volume 152 (1992)* 269

*Notes to authors* 271